arXiv:1108.1581v2 [math.NA] 9 Aug 2011

A Geometric Construction for the Evaluation of Mean Curvature

Pavel Grinfeld

7/20/2011

Abstract

We give a relationship that yields an effective geometric way of evaluating mean curvature of surfaces. The approach is reminiscent of the Gauss's contour based evaluation of intrinsic curvature. The presented formula may have a number of potential applications including estimating the normal vector and mean curvature on triangulated surfaces.

Given how brief is its derivation, it is truly surprising that this formula does not appear in the existing literature on differential geometry – at least according to the author's search. We hope to learn about a reference containing this result.

The presented formula admits a number of straightforward generalizations including to higher dimensions and submanifolds of co-dimension other than one. Here, we discuss the formula for a two dimensional surface embedded in a three dimensional Euclidean space.

Consider a regular patch P within a smooth surface S. Suppose that the area of P is A and that its smooth contour boundary is Γ. Let $\mathbf{N}$ be the normal to the surface and $\mathbf{n}$ be the exterior normal to Γ within S. Let B^α_α denote the mean curvature of S consistent with the direction of $\mathbf{N}$ **[1], [2]**. Then

$$\int_P \mathbf{N} B^\alpha_\alpha dS = \int_\Gamma \mathbf{n} d\Gamma. \tag{1}$$

In particular, for a patch shrinking to a point, we have

$$\mathbf{N} B^\alpha_\alpha = \lim \frac{1}{A} \int_\Gamma \mathbf{n} d\Gamma. \tag{2}$$

The right hand side of equation (2) is of coordinate free form and yields a pure geometric interpretation of mean curvature.

One immediate corollary of equation (1) is that the integral $\int_\Gamma \mathbf{n} d\Gamma$ vanishes for any contour in the interior of a minimal surface.

The derivation is quite brief and we cannot help but wonder whether equation (1) could have really been overlooked. The quantity $\mathbf{N} B^\alpha_\alpha$ equals the surface

divergence of the surface covariant basis $\mathbf{S}_\alpha$ [2, equivalent to eqn. (22) on p. 200]:

$$\mathbf{N}B^\alpha_\alpha = \nabla^\alpha \mathbf{S}_\alpha. \tag{3}$$

By Gauss's theorem on the patch P, we have

$$\int_P \nabla^\alpha \mathbf{S}_\alpha dS = \int_\Gamma n^\alpha \mathbf{S}_\alpha d\Gamma. \tag{4}$$

Recognizing that $n^\alpha \mathbf{S}_\alpha = \mathbf{n}$, we arrive at formula (1).

Formula (1) suggests a natural way of carrying over the concepts of the normal vector and mean curvature to triangulated surfaces. We note in advance that the proposed calculation yields an object analogous to the *product* $\mathbf{N}B^\alpha_\alpha$. Therefore, for minimal surfaces ($\mathbf{N}B^\alpha_\alpha = \mathbf{0}$) it will not yield the normal direction. Practical computational problems are almost sure to arise when computing normals for *nearly* minimal surfaces. On the other hand, the product $\mathbf{N}B^\alpha_\alpha$ is a more fundamental object than either $\mathbf{N}$ or B^α_α, and, more often than not, B^α_α arises precisely in the combination $\mathbf{N}B^\alpha_\alpha$.

Suppose that on a triangulated surface, triangles T_i, typically six in number, with areas A_i meet at the node O. Let $\mathbf{n}_i$ be unit vectors in the plane of T_i perpendicular to the triangle edges opposite of O and let $\mathbf{n}_i$ point away from O. Then the vector quantity

$$\mathbf{B} = \frac{\sum a_i \mathbf{n}_i}{\sum A_i} \tag{5}$$

may be taken as the definition of *vector mean curvature* for triangulated surfaces. The definition (5) is easily extended to the surface Laplacian of general invariant fields by replacing $\mathbf{n}_i$ with the normal derivative of the field.

The quantity $\mathbf{B}$ has a number of properties in common with its continuous analogue $\mathbf{N}B^\alpha_\alpha$, including:

i) $\mathbf{B} = \mathbf{0}$ for flat surfaces.

ii) More generally, $\mathbf{B}$ is the gradient of the total area with respect to the location of the vertices. Therefore, $\mathbf{B} = \mathbf{0}$ for minimal triangulated surfaces.

In conclusion, the central relationships (1) and (2) are analogous to Gauss's geometric construction for intrinsic curvature and may lead to effective computational applications. Surprisingly, we have not been able to locate these relationships in the existing literature and wish to share these useful identities as we continue our search.

References

[1] T. Levi-Civita. *The Absolute Differential Calculus (Calculus of Tensors).* Dover Publications, 1977.

[2] A. McConnell. *Applications of Tensor Analysis.* Dover Publications, New York, 1957.